

# Chapter 1

## Mathematical Preliminaries and Error Analysis

Per-Olof Persson  
persson@berkeley.edu

Department of Mathematics  
University of California, Berkeley

Math 128A Numerical Analysis

## Limits and Continuity

### Definition

A function  $f$  defined on a set  $X$  of real numbers has the *limit*  $L$  at  $x_0$ , written  $\lim_{x \rightarrow x_0} f(x) = L$ , if, given any real number  $\varepsilon > 0$ , there exists a real number  $\delta > 0$  such that

$$|f(x) - L| < \varepsilon, \quad \text{whenever } x \in X \text{ and } 0 < |x - x_0| < \delta.$$

### Definition

Let  $f$  be a function defined on a set  $X$  of real numbers and  $x_0 \in X$ . Then  $f$  is *continuous* at  $x_0$  if

$$\lim_{x \rightarrow x_0} f(x) = f(x_0).$$

The function  $f$  is continuous on the set  $X$  if it is continuous at each number in  $X$ .

## Limits of Sequences

### Definition

Let  $\{x_n\}_{n=1}^{\infty}$  be an infinite sequence of real or complex numbers. The sequence  $\{x_n\}_{n=1}^{\infty}$  has the *limit*  $x$  if, for any  $\varepsilon > 0$ , there exists a positive integer  $N(\varepsilon)$  such that  $|x_n - x| < \varepsilon$ , whenever  $n > N(\varepsilon)$ . The notation

$$\lim_{n \rightarrow \infty} x_n = x, \text{ or } x_n \rightarrow x \text{ as } n \rightarrow \infty,$$

means that the sequence  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

### Theorem

If  $f$  is a function defined on a set  $X$  of real numbers and  $x_0 \in X$ , then the following statements are equivalent:

- 1  $f$  is continuous at  $x_0$ ;
- 2 If the sequence  $\{x_n\}_{n=1}^{\infty}$  in  $X$  converges to  $x_0$ , then  $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$ .

## Derivatives

### Definition

Let  $f$  be a function defined in an open interval containing  $x_0$ . The function  $f$  is *differentiable* at  $x_0$  if

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

exists. The number  $f'(x_0)$  is called the *derivative* of  $f$  at  $x_0$ . A function that has a derivative at each number in a set  $X$  is *differentiable* on  $X$ .

### Theorem

If the function  $f$  is differentiable at  $x_0$ , then  $f$  is continuous at  $x_0$ .

## Derivative Theorems

### Theorem Rolle's Theorem

Suppose  $f \in C[a, b]$  and  $f$  is differentiable on  $(a, b)$ . If  $f(a) = f(b)$ , then a number  $c$  in  $(a, b)$  exists with  $f'(c) = 0$ .

### Theorem Mean Value Theorem

If  $f \in C[a, b]$  and  $f$  is differentiable on  $(a, b)$ , then a number  $c$  in  $(a, b)$  exists with

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

### Theorem Extreme Value Theorem

If  $f \in C[a, b]$ , then  $c_1, c_2 \in [a, b]$  exist with  $f(c_1) \leq f(x) \leq f(c_2)$ , for all  $x \in [a, b]$ . In addition, if  $f$  is differentiable on  $(a, b)$ , then the numbers  $c_1$  and  $c_2$  occur either at the endpoints of  $[a, b]$  or where  $f'$  is zero.

## Integrals

### Definition

The *Riemann integral* of the function  $f$  on the interval  $[a, b]$  is the following limit, provided it exists:

$$\int_a^b f(x) dx = \lim_{\max \Delta x_i \rightarrow 0} \sum_{i=1}^n f(z_i) \Delta x_i,$$

where the numbers  $x_0, x_1, \dots, x_n$  satisfy  $a = x_0 \leq x_1 \leq \dots \leq x_n = b$ , and where  $\Delta x_i = x_i - x_{i-1}$ , for each  $i = 1, 2, \dots, n$ , and  $z_i$  is arbitrarily chosen in the interval  $[x_{i-1}, x_i]$ .



## Floating Point Operations

### Finite-Digit Arithmetic

- Machine addition, subtraction, multiplication, and division:

$$x \oplus y = fl(fl(x) + fl(y)), \quad x \otimes y = fl(fl(x) \times fl(y))$$

$$x \ominus y = fl(fl(x) - fl(y)), \quad x \oslash y = fl(fl(x) \div fl(y))$$

- "Round input, perform exact arithmetic, round the result"

### Cancelation

- Common problem: Subtraction of nearly equal numbers:

$$fl(x) = 0.d_1d_2 \dots d_p\alpha_{p+1}\alpha_{p+2} \dots \alpha_k \times 10^n$$

$$fl(y) = 0.d_1d_2 \dots d_p\beta_{p+1}\beta_{p+2} \dots \beta_k \times 10^n$$

gives fewer digits of significance:

$$fl(fl(x) - fl(y)) = 0.\sigma_{p+1}\sigma_{p+2} \dots \sigma_k \times 10^{n-p}$$

## Rate of Convergence (Sequences)

### Definition

Suppose  $\{\beta_n\}_{n=1}^{\infty}$  is a sequence converging to zero, and  $\{\alpha_n\}_{n=1}^{\infty}$  converges to a number  $\alpha$ . If a positive constant  $K$  exists with

$$|\alpha_n - \alpha| \leq K|\beta_n|, \quad \text{for large } n,$$

then we say that  $\{\alpha_n\}_{n=1}^{\infty}$  converges to  $\alpha$  with *rate of convergence*  $O(\beta_n)$ , indicated by  $\alpha_n = \alpha + O(\beta_n)$ .

### Polynomial rate of convergence

- Normally we will use

$$\beta_n = \frac{1}{n^p},$$

and look for the largest value  $p > 0$  such that  $\alpha_n = \alpha + O(1/n^p)$ .

## Error Growth and Stability

### Definition

Suppose  $E_0 > 0$  is an initial error, and  $E_n$  is the error after  $n$  operations.

- $E_n \approx CnE_0$ : linear growth of error
- $E_n \approx C^n E_0$ : exponential growth of error

### Stability

- Stable* algorithm: Small changes in the initial data produce small changes in the final result
- Unstable* or *conditionally stable* algorithm: Large errors in final result for all or some initial data with small errors

## Rate of Convergence (Functions)

### Definition

Support that  $\lim_{h \rightarrow 0} G(h) = 0$  and  $\lim_{h \rightarrow 0} F(h) = L$ . If a positive constant  $K$  exists with

$$|F(h) - L| \leq K|G(h)|, \quad \text{for sufficiently small } h,$$

then we write  $F(h) = L + O(G(h))$ .

### Polynomial rate of convergence

- Normally we will use

$$G(h) = h^p,$$

and look for the largest value  $p > 0$  such that  $F(h) = L + O(h^p)$ .